

INCOMPLETE SELF-SIMILAR BLOW-UP IN A SEMILINEAR FOURTH-ORDER REACTION-DIFFUSION EQUATION

V.A. GALAKTIONOV

ABSTRACT. Blow-up behaviour for the 4th-order semilinear reaction-diffusion equation

$$(0.1) \quad u_t = -u_{xxxx} + |u|^{p-1}u \quad \text{in } \mathbb{R} \times \mathbb{R}_+, \quad p > 1,$$

is studied. For the classic semilinear heat equation from combustion theory

$$(0.2) \quad u_t = u_{xx} + u^p \quad (u \geq 0),$$

various blow-up patterns were investigated since 1970s, while the case of higher-order diffusion was studied much less. Blow-up self-similar solutions of (0.1),

$$u_-(x, t) = (T - t)^{-\frac{1}{p-1}} f(y), \quad y = \frac{x}{(T-t)^{1/4}},$$

are shown to admit global extensions for $t > T$ in an analogous similarity form:

$$u_+(x, t) = (t - T)^{-\frac{1}{p-1}} F(y), \quad y = \frac{x}{(t-T)^{1/4}}.$$

The continuity at $t = T$ is preserved in the sense that

$$u_-(x, T^-) = u_+(x, T^+) = C_1 |x|^{-\frac{4}{p-1}} \quad \text{for all } x \in \mathbb{R} \setminus \{0\} \quad (C_1 = \text{const.} \neq 0).$$

This is in a striking difference with blow-up for (0.2), which is known to be always *complete* in the sense that the minimal (proper) extension beyond blow-up is $u(x, t) \equiv +\infty$ for $t > T$. Difficult 4th-order dynamical systems for f and F are studied by a combination of various analytic, formal, and careful numerical methods. Other non-similarity patterns for (0.1) with non-generic *complete blow-up* are also discussed.

1. INTRODUCTION: SELF-SIMILAR BLOW-UP PATTERNS OF HIGHER-ORDER REACTION-DIFFUSION EQUATIONS AND MAIN RESULTS

1.1. Fourth-order RDE and blow-up. This paper continues the study began in [5, 13, 18, 25] of blow-up patterns for the *fourth-order reaction-diffusion equation* (the RDE-4)

$$(1.1) \quad u_t = -u_{xxxx} + |u|^{p-1}u \quad \text{in } \mathbb{R} \times \mathbb{R}_+, \quad \text{where } p > 1.$$

For applications of such higher-diffusion models, see surveys and references in [5, 25]. In general, higher-order semilinear parabolic equations arise in many physical applications such as thin film theory, convection-explosion theory, lubrication theory, flame and wave

Date: February 6, 2009.

1991 Mathematics Subject Classification. 35K55, 35K40 .

Key words and phrases. 4th-order semilinear parabolic equation, incomplete blow-up, self-similar solutions, similarity extensions.

propagation (the Kuramoto-Sivashinsky equation and the extended Fisher-Kolmogorov equation), phase transition at critical Lifschitz points, bi-stable systems and applications to structural mechanics; the effect of fourth-order terms on self-focusing problems in nonlinear optics are also well-known in applied and mathematical literature. For a systematic treatment of extended KPPF-equations; see in Peletier–Troy [36].

The RDE–4 (1.1) has the bi-harmonic diffusion operator $-D_x^4$ and is a higher-order counterpart of classic semilinear heat equation from combustion theory [41]

$$(1.2) \quad u_t = u_{xx} + |u|^{p-1}u \quad \text{in } \mathbb{R} \times \mathbb{R}_+,$$

with many well-known properties including various aspects of blow-up behaviour; see a number of well-known monographs [4, 38, 22, 34, 33, 24, 16, 37]. Surveys in [5] and in a more recent paper [18] contain necessary information concerning relations between these models (1.1) and (1.2) and description of similarity and other blow-up patterns for (1.1).

Note that another related fourth-order one-dimensional semilinear parabolic equation

$$(1.3) \quad u_t = -u_{xxxx} - [(2 - (u_x)^2)u_x]_x - \alpha u + qe^{su},$$

where α , q and s are positive constants obtained from physical parameters, occurs in the Semenov-Rayleigh-Benard problem [30], where the equation is derived in studying the interaction between natural convection and the explosion of an exothermically-reacting fluid confined between two isothermal horizontal plates. This is an evolution equation for the temperature fluctuations in the presence of natural convection, wall losses and chemistry. It can be considered as a formal combination of the equation derived in [27] (see also [6]) for the Rayleigh-Benard problem and of the Semenov-like energy balance [39, 12] showing that natural convection and the explosion mechanism may reinforce each other; see more details on physics and mathematics of blow-up in [25]. In a special limit, (1.3) reduces to the *generalized Frank-Kamenetskii equation*

$$(1.4) \quad u_t = -u_{xxxx} + e^u,$$

which plays a role of a natural extension of the classic Frank-Kamenetskii equation

$$(1.5) \quad u_t = \Delta u + e^u,$$

derived in solid fuel theory in the 1930s, [11].

Thus, similar to the second-order equations pair (1.2), (1.5), where both have an equal physical significance, we choose the RDE–4 (1.1) as a leading model, though some of the results are naturally applied to (1.4). Note that (1.1) can be considered as a non-mass-conservative counterpart of the well-known *limit unstable Cahn–Hilliard equation* from phase transition,

$$(1.6) \quad u_t = -u_{xxxx} - (|u|^{p-1}u)_{xx} \quad \text{in } \mathbb{R} \times \mathbb{R}_+,$$

which is known to admit various families of blow-up solutions; see [10] for a long list of references. It is key that the mass-conservation for L^1 -solutions of (1.6) naturally demands existence of an extension after blow-up, which was studied in [17]. This is not that

straightforward for the current model (1.1). Indeed, (1.1) is related to the famous mass-conserving divergence *Kuramoto–Sivashinsky equation* from flame propagation theory

$$(1.7) \quad u_t = -u_{xxxx} - u_{xx} + uu_x \quad \text{in } \mathbb{R} \times \mathbb{R}_+,$$

which always admits global solutions, so no blow-up for (1.7) exists.

As usual in PDE theory, *blow-up* means that, in the Cauchy problem (the CP) for (1.1) (or (1.2)) with bounded smooth initial function $u_0(x)$, the classic bounded solution $u = u(x, t)$ exists in $\mathbb{R} \times (0, T)$, while

$$(1.8) \quad \sup_{x \in \mathbb{R}^N} |u(x, t)| \rightarrow +\infty \quad \text{as } t \rightarrow T^-,$$

where $T \in \mathbb{R}_+ = (0, +\infty)$, depending on given data u_0 , is called the *blow-up time* of the solution $u(x, t)$. It is also convenient to use the quite popular nowadays auxiliary classification from Hamilton [29], where Type I blow-up means the solutions satisfying, for some constant $C > 0$ (depending on u),

$$(1.9) \quad \begin{aligned} \text{Type I: } & (T - t)^{\frac{1}{p-1}} |u(x, t)| \leq C \quad \text{as } t \rightarrow T^-, \text{ and, otherwise} \\ \text{Type II: } & \limsup_{t \rightarrow T^-} (T - t)^{\frac{1}{p-1}} \sup_x |u(x, t)| = +\infty \end{aligned}$$

(Type II also called *slow* blow-up in [29]). In R–D theory, blow-up with the first dimensional estimate in (1.9) was usually called of *self-similar* rate, while Type II was referred to as *fast* and non self-similar; see [24] and [38].

1.2. Main results: on incomplete blow-up and extended global semigroup. Unlike the previous papers, we are now more interested in a possible extension of blow-up solutions beyond blow-up, i.e., for $t > T$. For (1.2), this is not possible: blow-up of any its solution $u \geq 0$ is known to be *complete*, i.e., the proper (minimal) extension is

$$(1.10) \quad u(x, t) = +\infty \quad \text{in } \mathbb{R} \times (T, +\infty).$$

In fact, (1.10) is guaranteed by the Maximum Principle; see [23] and [16, Ch. 6,7] for main concepts and results of extended semigroup theory and further references.

Our main goal is to justify, that, rather surprisingly for us, for the RDE–4 (1.1),

$$(1.11) \quad \boxed{\text{self-similar blow-up is } \textit{incomplete} \text{ and admits self-similar extension for } t > T.}$$

Actually, (1.11) well corresponds to *Leray’s scenario* of self-similar blow-up and similarity extension beyond, proposed by Jean Leray in 1934 for the Navier–Stokes equations in $\mathbb{R}^3$; see [31, p. 245] for the precise formulation and [17] for recent discussions. For equation (1.1), (1.11) has an important corollary:

$$(1.12) \quad \boxed{\text{the number of self-similar extensions for } t > T \text{ is not more than countable}}$$

(actually, it is finite, but a proof exists in the analytic case $p = 3, 5 \dots$ only). This confirms a plausible existence of an extended semigroup of unique global proper (“minimal”) solutions of the Cauchy problem for (1.1).

We will also present some convincing facts that self-similar blow-up is a generic (structurally stable) one for (1.1), so that (1.11) implies a certain possibility of existence of a kind of extended semigroup theory of unique global proper solutions of (1.1) defined for

all $t > 0$. For (1.2), this is well known; see [23] as a guide. For (1.1), this question will be also discussed but essentially remains open.

For (1.2), the property (1.11) is revealed for the RDE-2 in $\mathbb{R}^N$ only, i.e., for the equation

$$(1.13) \quad u_t = \Delta u + u^p, \quad \text{where } p > p_{\text{Sob}} = \frac{N+2}{N-2}, \quad N > 2;$$

see [23]. However, even for (1.13), such an incomplete blow-up is not generic in the sense that, for almost all (a.a.) solutions, blow-up is expected to be complete. This is not fully justified rigorously, though the results imply this at least in the radial geometry.

Thus, according to (1.11), the RDE-4 (1.1) exhibits more flexibility in adapting blow-up solutions for $t > T$, though the analysis becomes essentially more difficult. To justify (1.11) in Section 2 (blow-up patterns for $t < T$) and 3 (global patterns for $t > T$), we will use a variety of methods including some analytical ones, but the final conclusions will eventually depend on careful numerical experiments. This is an unavoidable feature of the study of the 4th-order dynamical systems to be derived, which seems do not admit a fully mathematically rigorous investigation. As a key issue, we claim that

$$(1.14) \quad \boxed{\text{incomplete self-similar blow-up (1.11) for (1.1) has a pure dimensional nature.}}$$

Note that earlier, by a similar reason (1.14), existence of extended solutions was obtained [17] for the Cahn–Hilliard equation (1.6) for $p = 3$:

$$(1.15) \quad u_t = -u_{xxxx} - (u^3)_{xx} \quad \text{in } \mathbb{R} \times \mathbb{R}_+,$$

for which existence of a countable family of positive blow-up similarity solutions was established in [10]. However, since (1.15) is divergent and hence conservative by preserving the total mass of L^1 -solutions, existence of solutions beyond blow-up is more natural than for the non-conservative and non-divergent model (1.1).

In Section 4, we present other blow-up patterns for (1.1), introduced first in [13], which do not admit global extensions beyond blow-up, so belong to the case of *complete blow-up*. Fortunately, such complete blow-up is most plausibly non-generic for (1.1).

We hope that the conclusion (1.14) and others will be helpful for understanding singularity formation and extension concepts for $2m$ th-order nonlinear evolution PDEs including

$$(1.16) \quad u_t = -(-\Delta)^m u + |u|^{p-1}u \quad (m \geq 2),$$

which are definitely short of new ideas concerning construction of extended semigroups of unique global solutions. Note that some previous results in [5, 13] were already oriented to arbitrary $m = 2, 3, \dots$, though the present case of the even $m = 2$ in (1.1) will provide us with some surprises, especially for centre manifold patterns. In fact, dealing with (1.11), we show how to extend a blow-up solution beyond singularity towards hence guaranteeing existence of a *unique* (this is most desirable, but not easy to prove) continuation.

2. SELF-SIMILAR BLOW-UP PATTERNS

2.1. Blow-up similarity solutions. For convenience, we reduce the blow-up time to

$$T = 0,$$

simply meaning that, by shifting in time, the Cauchy problem for (1.1) is considered in, say, $(-1, +\infty) \times \mathbb{R}$ with initial data posed at $t = -1$.

Indeed, similarity blow-up is the simplest and most natural one for scaling invariant equations such as (1.1), where the behaviour as $t \rightarrow 0^-$ is given by a self-similar solution:

$$(2.1) \quad u_-(x, t) = (-t)^{-\frac{1}{p-1}} f(y), \quad y = \frac{x}{(-t)^{1/4}},$$

where $f(y) \not\equiv 0$ is a solution of the ODE:

$$(2.2) \quad \mathbf{A}_-(f) \equiv -f^{(4)} - \frac{1}{4} y f' - \frac{1}{p-1} f + |f|^{p-1} f = 0 \quad \text{in } \mathbb{R}, \quad f(\pm\infty) = 0.$$

We recall that, for (1.2), such nontrivial self-similar Type I blow-up (2.1),

$$(2.3) \quad u_-(x, t) = (-t)^{-\frac{1}{p-1}} f(y), \quad y = \frac{x}{(-t)^{1/2}} \implies f'' - \frac{1}{2} y f' - \frac{1}{p-1} f + |f|^{p-1} f = 0,$$

is nonexistent. This was first proved in Ad'jutov–Lepin in 1984 [1] (see [20] for first applications of the nonexistence to blow-up evolution). For (1.13), nonexistence in the subcritical Sobolev range $p \leq \frac{N+2}{N-2}$ was proved in Giga–Kohn [28] in 1985.

But this is not the case for the RDE-4 (1.1). Note that (2.2) is a difficult ordinary differential equation (ODE) with the non-coercive, non-monotone, and non-potential operators, so the problem is not variational in any weighted L^2 -spaces.

In what follows, by $f_0(y)$ we will denote the first monotone symmetric blow-up profile. We also deal with the second symmetric profile $f_1(y)$, which seems to be unstable, or, at least, less stable than f_0 .

The ODE (2.2) was studied in [5] by a number of analytic-branching and numerical methods. It was shown that (2.2) admits at least two different blow-up profiles with an algebraic decay at infinity. We also refer to [25, § 3] for further centre manifold-type arguments supporting this multiplicity result in a similar 4th-order blow-up problem. Without going into detail of such a study, we present a few illustrations only (mostly taken from [18]) and will address the essential dependence of similarity profiles $f(y)$ on p . In Figure 1, we present those pairs of solutions of (2.2) for $p = \frac{3}{2}$ and $p = 2$. All the profiles are symmetric (even), so satisfy the symmetry condition

$$(2.4) \quad f'(0) = f'''(0) = 0.$$

No non-symmetric blow-up was detected in numerical experiments (though there is no proof that such ones are nonexistent: recall that “moving plane” and Aleksandrov’s Reflection Principle methods do not apply to (1.1) without the MP). Figure 2 shows similar two blow-up profiles for $p = 5$.

2.2. Dimension of the “good” asymptotic bundle at infinity. To explain the nature of difficulties in proving existence of solutions of (2.2), let us describe the admissible behaviour for $y \gg 1$. It is not difficult to show that there exists a 2D bundle of such “good” asymptotics (see details in [9, § 3.3]): as $y \rightarrow +\infty$,

$$(2.5) \quad f(y) = [C_1 y^{-\frac{4}{p-1}} + \dots] + [C_2 y^{-\frac{2}{3}(\frac{p-3}{p-1})} e^{-a_0 y^{4/3}} + \dots], \quad \text{where } a_0 = 3 \cdot 2^{-\frac{8}{3}}$$




FIGURE 1. Two self-similar blow-up solutions of (2.2): $p = 1.5$ (a) and $p = 2$ (b).




FIGURE 2. Two self-similar blow-up solutions of (2.2) for $p = 5$.

and C_1 and C_2 are arbitrary parameters. The same symmetric bundle exists as $y \rightarrow \infty$. The first square bracket with algebraic decay is connected with the dominated linear terms:

$$(2.6) \quad -\frac{1}{4} f' y - \frac{1}{p-1} f + \dots = 0 \quad \implies \quad f(y) = C_1 y^{-\frac{4}{p-1}} + \dots$$

In the second square bracket in (2.5), we see a typical WKBJ-type two-scale asymptotics in ODE theory. The constant a_0 in (2.5) is obtained by substituting into the principal

part of (2.2) the pure exponential term:

$$(2.7) \quad -f^{(4)} - \frac{1}{4} y f' + \dots = 0, \quad \text{with} \quad f(y) \sim e^{ay^{4/3}} \implies a^3 = -\frac{1}{4} \left(\frac{3}{4}\right)^3,$$

whence the unique (real) root $-a_0$ with a negative real part. We do not go into details of the asymptotic expansions here, since such calculus are well known in deriving optimal exponential estimates of the fundamental solutions of higher-order parabolic equations, which were first obtained in Evgrafov–Postnikov (1970) and Tintarev (1982); see Barbatis [2, 3] for key references and updated results.

Note also that (2.5) reminds a typical centre manifold structure of the origin $\{f = 0\}$ at $y = \infty$: the first term in (2.5) is a node bundle with algebraic decay, while the second one corresponds to “non-analytic” exponential bundle around any of algebraic curves.

Thus, a dimensionally well-posed shooting is characterized as follows:

$$(2.8) \quad \boxed{\textbf{Shooting:} \quad \text{using } \mathbf{2} \text{ parameters } C_{1,2} \text{ in (2.5) to satisfy } \mathbf{2} \text{ conditions (2.4).}}$$

In view of the analytic dependence of solutions of (2.2) on the parameters $C_{1,2}$ in the bundle (2.5)¹, the problem (2.2), (2.4) cannot have more than a countable set of solutions. Actually, the numerics and branching-homotopy approaches [5] confirm that in wide parameter ranges of $p > 1$ and $N \geq 1$, there exist not more than two solutions:

$$(2.9) \quad f_0(y) \text{ with } \{C_{10}(p), C_{20}(p)\}, \quad \text{and} \quad f_1(y) \text{ with } \{C_{11}(p), C_{21}(p)\}.$$

More precisely, existence of both the first (generic blow-up) profile $f_0(y)$ and the second one $f_1(y)$ are obtained by a “ μ -bifurcation” approach, when the ODE (2.2) is replaced by

$$(2.10) \quad -f^{(4)} - \mu y f' - \frac{1}{p-1} f + |f|^{p-1} f = 0,$$

with a parameter $\mu \geq 0$. Linearization about the constant equilibrium and y -scaling yield:

$$(2.11) \quad f = (p-1)^{-\frac{1}{p-1}} + Y, \quad y = \left(\frac{1}{4\mu}\right)^{\frac{1}{4}} z \implies \mathbf{B}^* Y + \frac{1}{4\mu} Y + O(Y^2) = 0,$$

where $\mathbf{B}^* = -D_z^4 - \frac{1}{4} z D_z$ is the well-known Hermite-type operator with the discrete spectrum $\sigma(\mathbf{B}^*) = \{-\frac{k}{4}, k = 0, 1, 2, \dots\}$ and eigenfunctions being generalized Hermite polynomials (4.6), [8]; see more details in Section 4.2. It then follows from classic bifurcation theory [40] that bifurcations occur in (2.11) when $-\frac{1}{4\mu}$ gets on the spectrum of $\mathbf{B}^*$, i.e.,

$$(2.12) \quad -\frac{1}{4\mu} = -\frac{k}{4} \implies \exists \text{ bifurcation points } \mu_k = \frac{1}{k} \quad \text{for } k = 2, 4, 6, \dots,$$

where by natural symmetry reasons we take into account even k ’s only. It turns out that the bifurcations at $\mu_2 = \frac{1}{2}$ and $\mu_4 = \frac{1}{4}$ are responsible for existence of f_0 and f_1 respectively; see [5, § 5-7] for details. For convenience, let us note that the first μ -branch of solutions of (2.10) originated at $\mu_2 = \frac{1}{2}$ (a subcritical bifurcation) is strictly monotone decreasing for $\mu \in (0, \frac{1}{2})$ giving the f_0 at $\mu = \frac{1}{4}$. The next μ -branch originated at $\mu_4 = \frac{1}{4}$ (a supercritical bifurcation) is not monotone and is increasing on some interval $\mu \in (\frac{1}{4}, \mu_*)$, so there is another non-zero element on it at $\mu = \frac{1}{4}$, which is precisely the f_1 . In general,

¹This is rather plausible via standard trends of ODE theory [7], but difficult to prove. For positive solutions, it is true, and is always straightforward for odd $p = 3, 5, \dots$, where the nonlinearity is analytic.

the existence and multiplicity study of solutions of (2.2) in [5, 18, 25] is difficult and tricky, so here and later on we will need to essentially rely on careful numerical evidence to check the actual matching of the flows.

The eventual similarity blow-up patterns (2.1) are characterized by their *final time profiles*: passing to the limit $t \rightarrow 0^-$ in (2.1) and using the expansion (2.5) yields

$$(2.13) \quad \text{if } C_1(p) \neq 0, \text{ then } u_-(x, t) \rightarrow C_1 |x|^{-\frac{4}{p-1}} \text{ as } t \rightarrow 0^-$$

uniformly on any compact subset of $\mathbb{R} \setminus \{0\}$.

2.3. $C_1 = 0$: final time profile as a measure. If $C_1 = 0$ in (2.5), i.e., $f(y)$ has exponential decay at infinity, then the limit in (2.13) is different and there appears a measure in the data $u_-(x, 0^-)$: in the sense of distributions,

$$(2.14) \quad C_1(p) = 0 : \quad |u_-(x, t)|^{\frac{p-1}{4}} \rightarrow E_- \delta(x), \quad t \rightarrow 0^-; \quad E_- = \int |f|^{\frac{p-1}{4}} < \infty.$$

Taking into account the sign of the solution, we have a measure in the “mass” sense:

$$(2.15) \quad C_1(p) = 0 : \quad |u_-(x, t)|^{\frac{p-5}{4}} u_-(x, t) \rightarrow e_- \delta(x), \quad t \rightarrow 0^-; \quad e_- = \int |f|^{\frac{p-5}{4}} f \neq 0.$$

It is difficult to prove analytically that (2.14) actually takes place at some $p = p_\delta > 1$, so numerical methods have been used [18] to support this idea. Namely, the following exponent, for which (2.14) holds, was detected:

$$(2.16) \quad p_\delta(1) = 1.40\dots, \quad \text{for which } E_- = 45.4244\dots$$

In Figure 3, we show such profile $f_0(y)$ in the case (2.16), accompanied by the second one $f_1(y)$, for which $C_1 \neq 0$.

2.4. On existence of similarity profiles: classification of blow-up and oscillatory bundles. We now prove existence of at least a single blow-up profile $f(y)$ satisfying (2.2), (2.4). We perform shooting from $y = +\infty$ by using the 2D bundle (2.5) to $y = 0$, where the symmetry condition (2.4) are posed (or to $y = -\infty$, where the same bundle (2.5) with $y \mapsto -y$ takes place). By $f = f(y; C_1, C_2)$, we denote the corresponding solution defined on some maximal interval

$$(2.17) \quad y \in (y_0, +\infty), \quad \text{where } y_0 = y_0(C_1, C_2) \geq -\infty.$$

If $y_0(C_1, C_2) = -\infty$, then the corresponding solution $f(y; C_1, C_2)$ is global and can represent a proper blow-up profile (but not often, see below). Otherwise:

$$(2.18) \quad y_0(C_1, C_2) > -\infty \implies f(y; C_1, C_2) \rightarrow \infty \text{ as } y \rightarrow y_0^+.$$

Note that “oscillatory blow-up” for the ODE close to $y = y_0^+$:

$$(2.19) \quad f^{(4)} = |f|^{p-1} f (1 + o(1)),$$

where $\limsup f(y) = +\infty$ and $\liminf f(y) = -\infty$ as $y \rightarrow y_0^+$, is nonexistent. The proof is easy and follows by multiplying (2.19) by f' and integrating between two extremum




FIGURE 3. Two self-similar blow-up solutions of (2.2) for $p = p_\delta = 1.40\dots$, when (2.14) holds for f_0 .

points (y_1, y_2) , where the former one y_1 is chosen to be sufficiently close to the blow-up value y_0^+ , whence the contradiction:

$$0 < \frac{1}{2} (f'')^2(y_1) \sim -\frac{1}{p+1} |f|^{p+1}(y_1) < 0.$$

We first study this set of blow-up solutions. These results are well understood for such fourth-order ODEs; see [26], so we omit some details.

Proposition 2.1. *The set of blow-up solutions (2.19) is four-dimensional.*

Proof. The first parameter is $y_0 \in \mathbb{R}$. Other are obtained from the principal part of the equation (2.19) describing blow-up via (2.2) as $y \rightarrow y_0^+$. We apply a standard perturbation argument to (2.19). Omitting the $o(1)$ -term and assuming that $f > 0$, we find its explicit solution

$$(2.20) \quad f_0(y) = A_0(y - y_0)^{-\frac{4}{p-1}}, \quad A_0^{p-1} = \Phi(-\frac{4}{p-1}), \quad \Phi(m) \equiv m(m-1)(m-2)(m-3).$$

For convenience, the graph of $\Phi(m)$ is shown in Figure 4. Note that it is symmetric relative to $m_0 = \frac{3}{2}$, at which $\Phi(m)$ has a local maximum:

$$(2.21) \quad \Phi(\frac{3}{2}) = \frac{9}{16}.$$

By linearization, $f = f_0 + Y$, we get Euler's ODE:

$$(2.22) \quad (y - y_0)^4 Y^{(4)} = p A_0^{p-1} \equiv p \Phi(-\frac{4}{p-1}).$$




FIGURE 4. The graph of function $\Phi(m)$ in (2.20), and towards solutions of (2.22).

It follows that the general solution is composed from the polynomial ones with the following characteristic equation:

$$(2.23) \quad Y(y) = (y - y_0)^m \implies \Phi(m) = p\Phi\left(-\frac{4}{p-1}\right).$$

Since the multiplier $p > 1$ in the last term in (2.23) and $m = -\frac{4}{p-1}$ is a solution if this “ $p \times$ ” is omitted, this algebraic equation for m admits a unique positive solution $m_+ > 3$, a negative one $m_- < -\frac{4}{p-1}$, which is not acceptable by (2.20), and two complex roots $m_{1,2}$ with $\text{Re } m_{1,2} = \frac{3}{2} > 0$. Therefore, the general solution of (2.19) about the blow-up one (2.20), for any fixed y_0 , has a 3D stable manifold. $\square$

Thus, according to Proposition 2.1, the blow-up behaviour with a fixed sign (2.18) (i.e., non-oscillatory) is generic for the ODE (2.2). However, this 4D blow-up bundle together with the 2D bundle of good solutions (2.5) as $y \rightarrow \pm\infty$ are not enough to justify the shooting procedure. Indeed, by a straightforward dimensional estimate, an extra bundle at infinity is missing.

To introduce this new oscillatory bundle, we begin with the simpler ODE (2.19), without the $o(1)$ -term, and present in Figure 5 the results of shooting of a “separatrix” that lies between orbits, which blow-up to $\pm\infty$. Obviously, this separatrix is a periodic solution of this equation with a potential operator. Such variational problems are known to admit periodic solutions of arbitrary period.

Thus, Figure 5 fixed a bounded oscillatory (periodic) solution as $y \rightarrow +\infty$. When we return to the original equation (2.2), which is not variational, we still are able to detect a more complicated oscillatory structures at $y = \infty$. Namely, these are generated by the principal terms in

$$(2.24) \quad f^{(4)} = -\frac{1}{4}f'y + |f|^{p-1}f + \dots \quad \text{as } y \rightarrow \infty.$$




FIGURE 5. A periodic solution of $f^{(4)} = |f|^{p-1}f$ as a separatrix: $p = 5$.

Similar to Figure 5, in Figure 6, we present the result of shooting (from $y = -\infty$, which is the same by symmetry) of such oscillatory solutions of (2.2) for $p = 5$. It is easy to see that such oscillatory solutions have increasing amplitude of their oscillations as $y \rightarrow \infty$, which, as above, is proved by multiplying (2.24) by f' and integrating over any interval y_1, y_2 between two extrema. Figure 7 shows shooting of similar oscillatory structures at infinity for $p = 7$ (a) and $p = 2$ (b). It is not very difficult to prove that the set of such oscillatory orbits at infinity is 1D and this well corresponds to the periodic one in Figure 5 depending on the single parameter being its arbitrary period.

By $C_2^\pm(C_1)$ in Figure 6, we denote the values of the second parameters C_2 such that, for a fixed $C_1 \in \mathbb{R}$, the solutions $f(y; C_1, C_2^\pm)$ blow up to $\pm\infty$ respectively. These values are necessary for shooting the symmetry conditions (2.4).

Thus, overall, using two parameters $C_{1,2}$ in the bundle (2.5) for $y \gg 1$ leads to a well-posed problem of a 2D–2D shooting:

$$(2.25) \quad \text{find } C_{1,2} \text{ such that: } \begin{cases} y_0(C_1, C_2) = -\infty, & \text{and} \\ \text{no oscillatory behaviour as } y \rightarrow -\infty. \end{cases}$$

Concerning the actual proof of existence via shooting of at least a single blow-up patterns $f_0(y)$, by construction and oscillatory property of the equation (2.2), we first claim that in view of continuity relative to the parameters,

$$(2.26) \quad \text{for any } C_1 > 0, \text{ there exists } C_2^*(C_1) \in (C_2^-(C_1), C_2^+(C_1)) \text{ such that } f'''(0) = 0.$$

We next change C_1 to prove that at this $C_2^*(C_1)$ the derivative $f'(0)$ also changes sign. Indeed, one can see that

$$(2.27) \quad f'(0; C_1, C_2^*(C_1)) > 0 \quad \text{for } C_1 \ll 1 \quad \text{and} \quad f'(0; C_1, C_2^*(C_1)) < 0 \quad \text{for } C_1 \gg 1.$$




FIGURE 6. Shooting an oscillatory solution at infinity of (2.2): $p = 5$.




FIGURE 7. Shooting an oscillatory solution at infinity of (2.2): $p = 7$ (a) and $p = 2$ (b).

Actually, this means for such essentially different values of C_1 , the solution $f(y; C_1, C_2^*(C_1))$ has first oscillatory “humps” for $y > 0$ and $y < 0$ respectively. By continuity in C_1 , (2.27) implies existence of a C_1^* such that

$$(2.28) \quad f'(0; C_1^*, C_2^*(C_1^*)) = 0,$$

which together with (2.26) induced the desired solution. Overall, the above geometric shooting well corresponds to that applied in the standard framework of classic ODE theory, so we do not treat this in greater detail. However, we must admit that proving analogously existence of the second solution $f_0(y)$ (detected earlier by not fully justified

arguments of homotopy and branching theory and confirmed numerically) is an open problem. A more difficult open problem is to show why the problem (2.25) does not admit non-symmetric (non-even) solutions $f(y)$ (or does it?).

3. SELF-SIMILAR EXTENSIONS BEYOND BLOW-UP

3.1. Global similarity solutions. Following [23] devoted to (1.13) and actually using Leray's blow-up scenario [31], we suppose that the simplest way of extending of self-similar blow-up patterns for $t > 0 = T$ is using again similarity global patterns:

$$(3.1) \quad u_+(x, t) = t^{-\frac{1}{p-1}} F(y), \quad y = \frac{x}{t^{1/4}},$$

where F is a solution of the ODE

$$(3.2) \quad \mathbf{A}_+(F) \equiv -F^{(4)} + \frac{1}{4} y F' + \frac{1}{p-1} F + |F|^{p-1} F = 0 \quad \text{in } \mathbb{R}, \quad F(\pm\infty) = 0.$$

In comparison with the blow-up one (2.2), in (2.2) two linear terms in the middle have changed their signs. We will show that this essentially changes the dimension of the asymptotic bundles and hence overall matching results. Note that *exponentially decaying* solutions of (3.2) were already studied in [19] (these are important for extensions of the blow-up of type (2.14)), but now we need other types of solutions with algebraic decay.

For convenience, we always impose the same symmetry conditions

$$(3.3) \quad F'(0) = F'''(0) = 0,$$

and by $F_0(y)$ we will denote those profiles, which can be considered for the role of an extension of the blow-up profiles f_0 . Then $F_0(y)$ is not necessarily monotone for $y > 0$ and even its uniqueness is rather questionable; see numerical analysis below.

3.2. Dimensional analysis of matching for $C_1 \neq 0$. We first apply to (3.2) a simple test such as (2.7) to get the dimension of the exponentially decaying bundle:

$$(3.4) \quad F(y) \sim e^{ay^{4/3}} \implies a^3 = \frac{1}{4} \left(\frac{3}{4}\right)^3,$$

where the algebraic equation for $a \in \mathbb{C}$ admits *two* roots with negative real parts:

$$(3.5) \quad a_{\pm} = a_0 \left(-\frac{1}{2} \pm i \frac{\sqrt{3}}{2}\right), \quad \text{where } a_0 = 3 \cdot 2^{-\frac{8}{3}}.$$

Therefore, instead of (2.5), the bundle of such asymptotic orbits is 3D: as $y \rightarrow +\infty$,

$$(3.6) \quad \begin{aligned} F(y) = & \left[C_1 y^{-\frac{4}{p-1}} + \dots \right] \\ & + \left\{ y^{-\frac{2}{3}(\frac{p+1}{p-1})} e^{-\frac{a_0}{2} y^{\frac{4}{3}}} \left[C_2 \cos\left(\frac{a_0 \sqrt{3}}{2} y^{\frac{4}{3}}\right) + C_3 \sin\left(\frac{a_0 \sqrt{3}}{2} y^{\frac{4}{3}}\right) + \dots \right] \right\}, \end{aligned}$$

where C_2 and C_3 are arbitrary parameters. We consider here the case $C_1 \neq 0$, and will treat the special one $C_1 = 0$, with the data-measure (2.14), later on.

The point is that the first constant $C_1 \neq 0$ in (3.6) is fixed by the blow-up limit (2.13):

$$(3.7) \quad u_+(x, 0^+) = C_1 |x|^{-\frac{4}{p-1}}.$$




FIGURE 8. Blow-up profile $f_0(y)$ and the corresponding global one $F_0(y)$ with the same $C_1 = 77.76\dots$ for $p = 2$.

Overall, (2.13) and (3.7) will provide us with the necessary continuity of the unbounded self-similar solution at the blow-up time in the present case with $C_1 \neq 0$:

$$(3.8) \quad \boxed{u_-(x, 0^-) = u_+(x, 0^+) \quad \text{for } x \neq 0, \text{ i.e., almost everywhere (a.e.) in } \mathbb{R}.}$$

Then $\{f_0, F_0\}$ is called a *global extension similarity pair*, or simply an *extension pair*.

Thus, we again arrive at a dimensionally well-posed shooting: for a fixed values $C_1 \neq 0$,

$$(3.9) \quad \boxed{\textbf{Shooting:} \quad \text{using } \mathbf{2} \text{ parameters } C_{2,3} \text{ in (3.6) to satisfy } \mathbf{2} \text{ conditions (3.3).}}$$

3.3. Numerical analysis of self-similar blow-up extension for $C_1 \neq 0$. However, the actual solvability of the problem lying behind (3.8) is difficult for a rigorous analytic study. The main point is that the solvability is very much C_1 -dependent (and mostly is nonexistent for large C_1), so in many cases we will need again to rely on careful numerics.

In Figure 8, we show positive results of shooting the pair $f_0(y)$ and $F_0(y)$ for $p = 5$. For comparison, we also put therein some other global profiles $F(y)$ corresponding to other values of C_1 in (3.6) from $C_1 = 1$ up to $C_1 = 90$, i.e., larger than the required $77.76\dots$. In the next Figure 9, we show the enlarged asymptotic tails of all those profiles.

Figure 10 shows the extension pair $\{f_0, F_0\}$ for $p = 5$, where $C_1 = 1.843\dots$. In Figure 11, for the same case of $p = 5$, we present extra global F -profiles for smaller $C_1 = 0.2$ and larger $C_1 = 2$ values. We should note that convergence for the global F -problem is very slow and we succeeded in getting a few reliable numerics only. In all the cases, we have used the `bvp4c` solver of the `MatLab` with the enhanced accuracy with tolerances up to

$$(3.10) \quad \text{Tols} \sim 10^{-10}.$$




FIGURE 9. Enlarged tails of blow-up profile $f_0(y)$ and the corresponding global one $F_0(y)$ from Figure 8 with the same $C_1 = 77.76\dots$ for $p = 2$.




FIGURE 10. Blow-up profile $f_0(y)$ and the corresponding global one $F_0(y)$ with the same $C_1 = 1.843\dots$ for $p = 5$.

Note that not all the similarity blow-up profiles are assumed to have a global extension. For instance, Figure 12 explains nonexistence of a global profile $F_1(y)$ for the second blow-up one $f_1(y)$ from Figure 2 for $p = 5$. We present here the results of non-converging for




FIGURE 11. The pair $\{f_0(y), F_0(y)\}$ with $C_1 = 1.843\dots$ for $p = 5$ plus extra F -profiles.




FIGURE 12. Towards nonexistence of F_1 for the second blow-up profile $f_1(y)$, with $C_1 = 4.446\dots$ for $p = 5$.

the equation (3.2) with a sufficiently large

$$C_1 = 4.446\dots$$

Recall that, for $C_1 = 2$ from Figure 11(b), such a profile $F_0(y)$ exists, but seems non-existent for larger values of C_1 . Possibly, this means that there exists a critical maximal value of C_1 determining the optimal upper bound for existence of F -profiles.

3.4. On extension of a measure for $C_1 = 0$. Consider the special case (2.16), where, as shown in (2.14),

$$(3.11) \quad p = p_\delta = 1.40\dots : \quad |u_-(x, 0^-)|^{\frac{p-1}{4}} = E_- \delta(x), \quad \text{where} \quad E_- = 45.4244\dots$$

The corresponding blow-up profile $f_0(y)$ has been already shown in Figure 3.

If $C_1 = 0$ in (2.5), one then needs to get the global extension profile $F_0(y)$ also with $C_1 = 0$ in (3.6). This problem was studied in [19] by a bifurcation-branching approach. It was shown that there exists a countable sequence of critical exponents

$$(3.12) \quad p_l = 1 + \frac{4}{1+l}, \quad l = 0, 1, 2, \dots,$$

such that in $\{p > p_l\}$ there exists a global p -branch of solutions $F_l(y)$ of (3.2), that can be extended unboundedly as $p \rightarrow +\infty$. The behaviour of the branches near bifurcation points (3.12) was proved to be:

$$(3.13) \quad F_{0l}(y) = \gamma_l(p - p_l)^{\frac{1+l}{4}} [\psi_l(y) + o(1)] \quad \text{as} \quad p \rightarrow p_l^+; \quad l = 0, 1, 2, \dots,$$

where $\gamma_l > 0$ are some constants and $\psi_l(y)$ are eigenfunctions (4.9) of the operator (4.7).

Before constructing the extension pair for the case (3.11), note that this p_δ , within our accuracy, is close to the spectrum (3.12), and, precisely,

$$(3.14) \quad p_9 = 1 + \frac{4}{10} = 1.40.$$

Note also that the critical exponents (3.12) are concentrated about $p = 1^+$ for large l , and, in view of the asymptotics (3.13), the p -branches are rather plain close to $p = p_\delta$. Overall, this shows that there exist many global profiles $F_{0l}(y)$ satisfying (3.11).

Figure 13 shows a typical global similarity profile $F_0(y)$, though by the branching (3.13), there exists a countable set of such solutions of (3.2) with exponential decay. Therefore, it is difficult to identify which p -branch this profile F_0 belongs to. The measure characteristic of this $F_0(y)$ in Figure 13 is:

$$(3.15) \quad E_+ = 13.2893\dots$$

The computations have been performed with the enhanced tolerances (3.10), since many profiles F_0 are small enough due to (3.13) for $p \approx p_\delta$. On the other hand, by the same branching, for $l \gg 1$ with p_l more closer to 1^+ , there exist F_0 's with arbitrarily large E_{+l} .

Indeed, we see that the value (3.15) for a global extension does not match the blow-up one in (3.11):

$$(3.16) \quad p = p_\delta = 1.40\dots : \quad E_- > E_+.$$

Recall that, due to (3.12), there also exist global profiles F_{0l} such that $E_- < E_{+l}$. However, all these discrepancies do not undermine a possibility of extension of the data (2.14) in a self-similar way given by (3.1). An effective scenario of such a matching was obtained and confirmed in [10] for the related limit unstable Cahn–Hilliard equation (1.15), which preserves the mass of the L^1 -solutions. Namely, it was shown [10, § 4] that the rescaled blow-up solution can converge in the local topology (4.12) to a self-similar profile with a different mass (L^1 -norm for positive solutions). In practice, this means that close to blow-up time (as $t \rightarrow 0^-$), negative humps are created that eventually disappear at infinity,




FIGURE 13. A typical global similarity profile $F_0(y)$ for $p = p_\delta = 1.40\dots$.

and hence do not violate the matching. This “mass drift” is performed in the rescaled variables as $t \rightarrow 0^-$, so does not mean existence of the actual mass transferring mechanism in the x -variable that might be unavailable in the PDE. There is no a rigorous proof.

We expect that a similar mechanism can be explored to neglect the mass “defect”, and, for $t > 0$, we may observe a global profile $F_{0l}(y)$, which minimizes the loss of the mass $E_- - E_{+l}$. The positive mass defect will then create two negative/positive humps that disappear at infinity in the rescaled variable y as $t \rightarrow 0^+$. In this mass interpretation, we have to use the measure-like data obtained in the limit (2.15). Recall that the whole set of global profiles $\{F_{0l}\}$ at $p = p_\delta$ is countable, so that this minimization makes sense. It looks then like a certain “mass-discontinuity” is assumed at $t = 0^+$, though this happens in the $y = x/t^{1/4}$ variable and hence the real discontinuity is not available in the x -one, which is prohibited for this parabolic flow.

There are several open problems here, and the further analysis of the actual extension pair $\{f_0, F_{0l}\}$ based on enhanced *PDE numerics* is quite necessary and desirable to clarify this difficult transition blow-up singularity phenomenon.

3.5. Towards extended semigroup theory. Thus, the above results, at least at a qualitative formal level, imply that self-similar blow-up for (1.1) (which is assumed to be generic and most structurally stable; see the next section) admits a proper extension beyond blow-up time. Moreover, we also can claim that, for $C_1 \neq 0$:

(3.17) there exists at most a finite number of extension pairs $\{f_0, F_0\}$.

For $C_1 = 0$, i.e., for $p = p_\delta$, due to the critical bifurcation exponents (3.12), we expect a countable sequences of the extension pairs, but the discrepancy of their “masses” at $t = 0$

could reinforce extra evolution mechanism of choosing the right ones. This remains an open problem.

Note that, in general, we cannot guarantee that a pair is unique. Nevertheless, in view of (3.17), even in the case of finite multiplicity of the profiles F_0 , there is a hope that

$$(3.18) \quad \text{there exists a “minimal” extension pair } \{f_0, \bar{F}_0\},$$

with $\bar{F}_0$ being “ordered” in a certain geometric-metric sense, e.g., F_0 with the same C_1 but with a most “less oscillatory” structure. Recall that f_0 is always monotone.

This *minimality property* can play a key role in an attempt (seems rather naive) to constructing of extended semigroup theory for blow-up solutions of (1.1) by using standard ideas of parabolic regularizations via smooth solutions $\{u_\varepsilon\}$ (cf. [23] for (1.2)),

$$(3.19) \quad u_\varepsilon : u_t = -u_{xxxx} + \varphi_\varepsilon(u), \quad \text{where} \quad \varphi_\varepsilon(u) = \frac{|u|^{p-1}u}{1+\varepsilon|u|^{p-1}u} \rightarrow |u|^{p-1}u \quad \text{as} \quad \varepsilon \rightarrow 0^+$$

uniformly on compact subsets. Since $\varphi_\varepsilon(u)$ is globally Lipschitz continuous, the CP for (3.19) with the same data u_0 has a unique global classical solution $u_\varepsilon(x, t)$. Then a proper (minimal) solution of (1.1) can be formally defined as a hypothetical limit

$$(3.20) \quad \bar{u}(x, t) = \lim_{\varepsilon \rightarrow 0} u_\varepsilon(x, t) \quad \text{in} \quad \mathbb{R} \times (-1, \infty),$$

where the limit is understood in a pointwise sense, since $u(x, t)$ is unbounded for $t \geq 0$. Moreover, no more restrictions on the topology of convergence in (3.20) can be imposed, since, for general solutions, the *blow-up set*:

$$(3.21) \quad B[u_0](t) = \{x \in \mathbb{R} : |u(x, t)| = \infty\},$$

has an unknown structure for $t \geq T$ (hopefully, of zero measure, which is not proved yet). Recall that even for much simpler nonlinear heat equations including (1.13), the limit extended semigroups are discontinuous in time in general; see [16, Ch. 7] for examples.

Proving existence of the limit (3.20) (even along a subsequence) and checking how it is related to the minimal $\bar{F}_0$ -extension of the blow-up self-similar solution (2.1) are difficult open problems (possibly, non-solvable in general). Anyway, it is indeed surprising that there is still a slight hope that extended semigroup theory of blow-up solutions for (1.1) can be at least partially developed along the lines of that for the second-order parabolic PDEs such as (1.2) [23, 16], where the Maximum Principle was always key.

3.6. On generalizations to $\mathbb{R}^N$. Finally, as a comment and an introduction to a future research, in Figure 14 we show the extension pair $\{f_0, F_0\}$ for the equation in $\mathbb{R}^N$,

$$(3.22) \quad u_t = -\Delta^2 u + |u|^{p-1}u,$$

in the case $p = 5$ for $N = 3$ (a) and 6 (b). Note the clear difference in the geometry of the corresponding global F_0 -profiles in (a) and (b). This is again a sign showing that the extension profiles F_0 can be nonunique.

The similarity solutions $u_\pm$ remain the same and the ODEs for radial patterns $\{f_0, F_0\}$ are easily obtained; see [18] for details on blow-up ones $u_-(x, t)$. The ODE problems for




FIGURE 14. The pair $\{f_0(y), F_0(y)\}$ for $p = 5$ and $N = 3$ (a) and $N = 6$ (b).

f_0 and F_0 in the radial geometry are as follows (now y stands for $|y| > 0$):

$$\begin{cases} -f^{(4)} - \frac{2(N-1)}{y} f''' - \frac{(N-1)(N-3)}{y^2} f'' + \frac{(N-1)(N-3)}{y^3} f' - \frac{1}{4} y f' - \frac{1}{p-1} f + |f|^{p-1} f = 0, \\ f'(0) = f'''(0) = 0, \quad f(y) = C_1 y^{-\frac{4}{p-1}} + \dots \quad \text{as } y \rightarrow \infty; \end{cases}$$


$$\begin{cases} -F^{(4)} - \frac{2(N-1)}{y} F''' - \frac{(N-1)(N-3)}{y^2} F'' + \frac{(N-1)(N-3)}{y^3} F' + \frac{1}{4} y F' + \frac{1}{p-1} F + |F|^{p-1} F = 0, \\ F'(0) = F'''(0) = 0, \quad f(y) = C_1 y^{-\frac{4}{p-1}} + \dots \quad \text{as } y \rightarrow \infty. \end{cases}$$

The asymptotic bundles (2.5) and (3.6) remain analogous to the case $N = 1$.


In Figure 15, we show the pairs $\{f_0, F_0\}$ again for $p = 5$ in dimensions $N = 8$ (a) and $N = 9$ (b). Finally, in Figure 16, the extension pair is shown for $p = 5$ and $N = 15$. Analogously to $N = 1$, the convergence for the global profile $F_0(y)$ is very slow, possibly in view of the multi-dimension of the bundle (3.6). However, for such profiles, we always observed a multiple convergence to such profiles from various initial data and/or varied accuracy and tolerances used in the **bvp4c** solver.

4. NON SELF-SIMILAR “LINEARIZED” PATTERNS: COMPLETE BLOW-UP

A countable set of such non-self-similar blow-up patterns for (1.1) were formally constructed in [13]; see also [18] for extra details and necessary historical comments. Below, we briefly describe some necessary new features of these patterns to reveal the main reasons for their complete blow-up. For the RDE-4, there is no hope to get a full rigorous justification of existence of such non-self-similar blow-up scenarios, so we feel free to perform a detailed formal construction. More advanced expansion and matching techniques in this direction can be found in [13] and also [14], where such a construction applied to non-singular absorption phenomena (regular flows with no blow-up), so a full mathematical justification is available therein.



(a) $N = 8$, $C_1 = 2.22\dots$



(b) $N = 9$, $C_1 = 3.38\dots$

FIGURE 15. The pair $\{f_0(y), F_0(y)\}$ for $p = 5$ and $N = 8$ (a) and $N = 9$ (b).




FIGURE 16. The pair $\{f_0, F_0\}$ for $p = 5$, $N = 15$.

4.1. Nonstationary rescaling. Dealing with non-self-similar blow-up, instead of (2.1), we use the full similarity scaling:

$$(4.1) \quad u(x, t) = (-t)^{-\frac{1}{p-1}} v(y, \tau), \quad y = \frac{x}{(-t)^{1/4}}, \quad \tau = -\ln(-t) \rightarrow +\infty \text{ as } t \rightarrow 0^-.$$

Then $v(y, \tau)$ solves the following parabolic equation:

$$(4.2) \quad v_\tau = \mathbf{A}_-(v) \equiv -v_{yyyy} - \frac{1}{4} y v_y - \frac{1}{p-1} v + |v|^{p-1} v \quad \text{in } \mathbb{R} \times \mathbb{R}_+,$$

where $\mathbf{A}_-$ is the stationary operator in (2.2), so that similarity profiles are simply stationary solutions of (4.2).

4.2. Linearization and spectral properties. Performing the standard linearization about the constant equilibrium in the equation (4.2) yields a perturbed equation:

$$(4.3) \quad v = f_* + Y, \quad f_* = (p-1)^{-\frac{1}{p-1}} \implies Y_\tau = (\mathbf{B}^* + I)Y + \mathbf{D}(Y),$$

where $\mathbf{D}(Y) = c_0 Y^2 + \dots$, $c_0 = \frac{p}{2}(p-1)^{\frac{1}{p-1}}$, is a quadratic perturbation as $Y \rightarrow 0$ and

$$(4.4) \quad \mathbf{B}^* = -D_y^4 - \frac{1}{4} y D_y \quad \text{in} \quad L_{\rho^*}^2(\mathbb{R}), \quad \rho^*(y) = e^{-a|y|^{4/3}}, \quad a \in (0, 3 \cdot 2^{-\frac{8}{3}}),$$

is the *adjoint Hermite operator* with some good spectral properties [8]:

Lemma 4.1. $\mathbf{B}^* : H_{\rho^*}^4(\mathbb{R}) \rightarrow L_{\rho^*}^2(\mathbb{R})$ is a bounded linear operator with the spectrum

$$(4.5) \quad \sigma(\mathbf{B}^*) = \{\lambda_k = -\frac{k}{4}, \quad k = 0, 1, 2, \dots\} \quad (= \sigma(\mathbf{B}), \quad \mathbf{B} = -D_y^4 + \frac{1}{4} y D_y + \frac{1}{4} I).$$

Eigenfunctions $\psi_k^(y)$ are k th-order generalized Hermite polynomials:*

$$(4.6) \quad \psi_k^*(y) = \frac{1}{\sqrt{k!}} [y^k + \sum_{j=1}^{\lfloor k/4 \rfloor} \frac{1}{j!} (D_y)^{2j} y^k], \quad k = 0, 1, 2, \dots,$$

and the subset $\{\psi_k^\}$ is complete in $L_{\rho^*}^2(\mathbb{R})$.*

As usual, if $\{\psi_k\}$ is the adjoint (Riesz) basis of eigenfunctions of the adjoint operator

$$(4.7) \quad \mathbf{B} = -D_y^4 + \frac{1}{4} y D_y + \frac{1}{4} I \quad \text{in} \quad L_\rho^2(\mathbb{R}), \quad \text{with} \quad \rho = \frac{1}{\rho^*},$$

with the same spectrum (4.5), the bi-orthonormality condition holds in $L^2(\mathbb{R})$:

$$(4.8) \quad \langle \psi_k, \psi_l^* \rangle = \delta_{kl} \quad \text{for any} \quad k, l.$$

The generating formula for ψ_k is as follows:

$$(4.9) \quad \psi_k(y) = \frac{(-1)^k}{\sqrt{k!}} F^{(k)}(y), \quad k = 0, 1, 2, \dots,$$

where $F(y)$ is the rescaled kernel of the fundamental solution $b(x, t)$ of $D_t + D_x^4$, i.e.,

$$(4.10) \quad b(x, t) = t^{-\frac{1}{4}} F(y), \quad y = \frac{x}{t^{1/4}}; \quad b_t = -b_{xxxx}, \quad b(x, 0) = \delta(x) \implies \mathbf{B}F = 0, \quad \int F = 1.$$

4.3. Why self-similar blow-up is expected to be generic. This can be connected with the spectral gaps in $\sigma(\mathbf{B}^* + I) = \{1 - \frac{k}{4}\}$ of the linearized operator in (4.3). Indeed, we observe here four positive unstable modes:

$$(4.11) \quad k = 0 \quad \text{and} \quad 1, \quad \text{and} \quad k = 2 \quad \text{and} \quad 3.$$

The first group is not taken into account since correspond to standard instabilities relative to perturbations of the time $T = 0$ and the space $x = 0$ of the blow-up point.

The crucial is the actual unstable mode with

$$(4.12) \quad k = 2, \quad \text{where} \quad \lambda_2 = \frac{1}{2} \quad \text{and} \quad \psi_2^*(y) = \frac{1}{\sqrt{2}} y^2.$$

We claim that this unstable mode ensures convergence to a non-trivial similarity stationary profile, which we used to denote by $f_0(y)$; see extra explanations in [25, § 3]. There is no still a proof of this intriguing fact on such a heteroclinic connection $f_* \rightarrow f_0$, but it

sounds rather reliably. Hence, the stable manifold of f_0 includes a part of the unstable manifold of the constant equilibrium $f_* = (p-1)^{-1/(p-1)}$, so that linearized patterns to be constructed must be less stable than f_0 , which then is expected to represent a most structurally stable (generic) blow-up pattern.

Note that the third unstable mode in (4.11) with $k = 3$ has nothing to do with the second similarity profile f_1 . Indeed, this mode corresponds to odd (anti-symmetric) perturbations, while $f_1(y)$ is an even function. The origin of existence of f_1 is more subtle and is explained in [5] by branching and homotopy theory. Moreover, according to [25, § 3.2], $f_1(y)$ is connected with the trivial equilibrium 0 via a semi-stable centre manifold behaviour to be presented below.

4.4. Inner expansion. Thus, in the *Inner Region* characterized by compact subsets in the similarity variable y , we assume a centre or a stable subspace behaviour as $\tau \rightarrow +\infty$ for the linearized operator $\mathbf{B}^* + I$ (see [14] for rigorous details in a simpler related problem):

$$(4.13) \quad \begin{aligned} \text{centre: } & Y(y, \tau) = a(\tau)\psi_4^*(y) + w^\perp \quad (\lambda_4 = -1, \ k = 4), \\ \text{stable: } & Y(y, \tau) = -Ce^{\lambda_k \tau}\psi_k^*(y) + w^\perp \quad (\lambda_k < -1, \ k > 4). \end{aligned}$$

As usual for the classic equation (1.2), we restrict to even values only,

$$(4.14) \quad k = 4, 6, 8, \dots,$$

since existence of patterns with odd k 's, i.e., having a non-symmetric blow-up structures is rather suspicious. However, for higher-order diffusion, such patterns cannot be excluded entirely, though this is not our business here.

For the centre subspace behaviour in (4.13), substituting the eigenfunctions expansion into equation (4.3) yields the following coefficient:

$$(4.15) \quad \dot{a} = a^2\gamma_0 + \dots, \quad \text{where} \quad \gamma_0 = c_0\langle(\psi_4^*)^2, \psi_4\rangle \implies a(\tau) = -\frac{1}{\gamma_0\tau} + \dots$$

It is crucial that (4.15) shows a clear *semi-stable* (saddle-node) structure of the equilibrium $a = 0$, which essentially depends on the sign of the coefficient γ_0 ; see below.

Note that for the matching purposes, one needs to assume that (see details in [13, 25]):

$$(4.16) \quad \text{if } \psi_k^*(0) > 0, \text{ then } \gamma_0 > 0 \quad \text{and} \quad C > 0.$$

Actually, for $k = 4$ it is calculated explicitly that

$$(4.17) \quad \gamma_0 = -c_0 136\sqrt{6} < 0,$$

so that the centre manifold patterns with the positive eigenfunction

$$(4.18) \quad \psi_4^*(y) = \frac{1}{\sqrt{24}}(y^4 + 24)$$

correspond to solutions that blow-up on finite interfaces, [25, § 3]. It seems reasonable that a full justification of such a behaviour can be done along the lines of classic invariant manifold theory (see e.g. [32]), though can be very difficult.

Overall, the whole variety of such asymptotics is characterized as follows:

$$(4.19) \quad Y(y, \tau) = a(\tau)y^k + \dots, \quad \text{where} \quad a(\tau) = - \begin{cases} \frac{1}{\gamma_0 \tau} + \dots & \text{for } k = 4, \\ Ce^{\lambda_k \tau} + \dots & \text{for } k > 4. \end{cases}$$

4.5. Outer region: matching. The asymptotics (4.13) is known [13] to admit matching with the *Outer Region*, being a Hamilton–Jacobi (H–J) one. More precisely, in the centre case with $k = 4$, according to (4.13), (4.15), we introduce the outer variable and obtain from (4.2) the following perturbed H–J equation:

$$(4.20) \quad \xi = \frac{y}{\tau^{1/4}} \implies v_\tau = -\frac{1}{4} \xi v_\xi - \frac{1}{p-1} v + |v|^{p-1} v + \frac{1}{\tau} \left(\frac{1}{4} \xi v_\xi - v_{\xi\xi\xi\xi} \right).$$

Passing to the limit as $\tau \rightarrow +\infty$ in such singularly perturbed PDEs is not easy at all even in the second-order case (see a number of various applications in [24]). Though currently not rigorously, we assume stabilization to the stationary solutions $f(\xi)$ satisfying the unperturbed H–J equation:

$$(4.21) \quad -\frac{1}{4} \xi f' - \frac{1}{p-1} f + |f|^{p-1} f = 0 \quad \text{in } \mathbb{R}.$$

This is solved via characteristics, where we have to choose the solution satisfying (4.19):

$$(4.22) \quad f(\xi) = (p-1)^{-\frac{1}{p-1}} - \frac{1}{\gamma_0} \xi^4 + \dots \quad \text{as } \xi \rightarrow 0.$$

Hence, integrating (4.21) with the condition (4.22) yields the unique H–J profile

$$(4.23) \quad f_0(\xi) = f_* (1 + c_* \xi^4)^{-\frac{1}{p-1}}, \quad \text{where} \quad c_* = \frac{1}{\gamma_0} (p-1)^{\frac{p}{p-1}}.$$

Note that since $\gamma_0 < 0$ according to (4.17), the profile (4.23) blows up as $c_* \xi^4 \rightarrow -1$; see analogous phenomena in [25]. For the $2m$ th-order PDE (1.16) with odd $m = 3, 5, \dots$, we have $\gamma_0 > 0$ and then (4.23) can represent standard blow-up patterns, [13].

Analogously, for the stable behaviour for $k > 4$ in (4.13), we use the following change:

$$(4.24) \quad \xi = e^{\frac{4-k}{4k} \tau} y \implies v_\tau = -\frac{1}{k} \xi v_\xi - \frac{1}{p-1} v + |v|^{p-1} v - e^{-\frac{k-4}{k} \tau} v_{\xi\xi\xi\xi}.$$

Passage to the limit $\tau \rightarrow +\infty$ and matching with Inner Region are analogous. The H–J profile is as follows:

$$(4.25) \quad f_C(\xi) = f_* (1 + c \xi^k)^{-\frac{1}{p-1}}, \quad \text{where} \quad c = C (p-1)^{\frac{p}{p-1}} > 0.$$

Overall, according to matching conditions (4.19) and (4.24), the whole set of such blow-up patterns consists of a countable set of different solutions for $k = 6, 8, \dots$ (as we have mentioned, the case $k = 4$ works for m odd only, and then $k = 2m$).

4.6. Final time profiles: the rout to complete blow-up. As a natural counterpart of the above asymptotic analysis, we introduce the *final time profiles* of solutions [13, § 4]:
CENTRE SUBSPACE PATTERNS: $k = 4$. This analysis is formal, since by (4.17) and (4.23) with $c_* < 0$ such patterns are not bounded. However, we keep the derivation formally assuming that case $\gamma_0 > 0$, which applies to equations (1.16) with odd $m = 3, 5, 7, \dots$,

[13]. Thus, marching of the outer region with the Outer Region II for $|x| > 0$ small, yields the following behaviour:

$$(4.26) \quad u(x, 0^-) = C_* |x|^{-\frac{4}{p-1}} |\ln |x||^{\frac{1}{p-1}} (1 + o(1)),$$

where $C_* > 0$ is a constant depending on the parameters p and γ_0 (assumed here to be positive) only and hence is independent of initial data u_0 . A more careful passing to the limit as $t \rightarrow 0^-$ in (4.1) by using the limit profile (4.23) shows that

$$(4.27) \quad C_* = f_* \left(\frac{4}{c_*} \right)^{\frac{1}{p-1}}.$$

STABLE SUBSPACE PATTERNS: $k > 4$. Such stable subspace patterns can be truly constructed for all $m = 2, 3, \dots$. Matching with the Outer Region II yields, as $x \rightarrow 0$,

$$(4.28) \quad u(x, 0^-) = C_* |x|^{-\frac{k}{p-1}} (1 + o(1)) \quad \text{for } k = 6, 8, \dots,$$

where constants C_* depend on initial function u_0 through the earlier constant C in (4.19).

Comparing the self-similar profile (2.13) with those in (4.26) and (4.28) shows the actual origin of *complete blow-up*: the later ones contain stronger singularities as initial data at $x = 0$ posed at $t = 0$. This is easier to see for the case $k = 4$ (actually, nonexistent by (4.17)), where the profiles differ by the unbounded factor

$$(4.29) \quad \sim |\ln |x||^{\frac{1}{p-1}} \rightarrow +\infty \quad \text{as } x \rightarrow 0.$$

Surely, the same and in a stronger manner happens for $k = 6, 8, \dots$.

Bearing in mind our self-similar blow-up patterns (2.1) with the asymptotics (2.5), the unbounded factor in (4.29) actually means that, in a certain natural but formal sense, we have to look for a similarity pattern with

$$(4.30) \quad C_1 = +\infty.$$

Moreover, for existence of a regular extension for $t > 0$, this would demand existence of a global patterns (3.1), (3.6) also with the coefficient (4.30). Of course, this is not possible that somehow reflects the nonexistence of a proper extension, meaning complete blow-up.

The actual proof of nonexistence of a regular solution of (1.1) with data (4.28) is not easy. This assumes proving nonexistence of a corresponding *very singular solution* for (1.1), which is a well-known issue for second-order semilinear and quasilinear heat equations. However, such results are also known for higher-order nonlinear parabolic equations with various nonlinearities, where the analysis without the Maximum Principle becomes essentially more involved. We refer to functional methods in [21], where necessary earlier references on the subject can be found.

5. FINAL REMARK: ON EVOLUTION COMPLETENESS OF BLOW-UP PATTERNS

After introducing and discussing all the blow-up patterns, we finally are in a position to announce another important aspect of our study. Namely, we claim that the whole set of self-similar and linearized patterns introduced above are *evolutionary complete* in a natural sense; see [15]. This purely means that the self-similar blow-up patterns (2.2)

together with all the linearized ones in Section 4 are expected to describe all possible types of blow-up that can occur for the PDE (1.1). We connect this claim with the standard completeness of the eigenfunctions (4.6) in $L^2_{\rho^*}$. However, since by (4.11) we have thrown away two unstable patterns for $k = 2$ and $k = 3$ (the modes with $k = 0$ and 1 do not occur in the evolution), one needs to replace these two ones. And this is done by including precisely *two* nonlinear self-similar patterns (2.1) with profiles f_0 and f_1 . Note that this may look irrelevant since the mode with $k = 3$ is spatially odd, while $f_1(y)$ was shown to be even. In fact, this is not that important, since we always mean a certain symmetrization of blow-up behaviour as $t \rightarrow T^-$ (no proof is still available), so that including two “nonlinear eigenfunctions” $f_{0,1}(y)$ is expected to be enough to govern non-trivial self-similar blow-up.

What is more crucial is that a standard centre manifold behaviour is prohibited by the sign in (4.17). Therefore, finally, according to evolution completeness arguments:

two similarity profiles $f_{0,1}(y)$ replace two modes, with $k = 2$ and 4.

Here we observe a usual “dimension preservation” by including into the countable set of linearized patterns (minus two modes for $k = 2, 4$) two nonlinear eigenfunctions $f_{0,1}(y)$.

Of course, another and obviously inevitable key aspect of the completeness speculation above is *nonexistence* of Type II blow-up in (1.9) for (1.1). Indeed, Type II blow-up patterns would destroy any completeness of the above functional set. Note that such Type II patterns were obtained for (3.22) for some large p in the supercritical Sobolev range

$$p \geq p_{\text{Sob}} = \frac{N+4}{N-4} \quad (N > 4);$$

see [18, § 5, 6] for details. We have some reliable evidence that such patterns are nonexistent in the subcritical range $p < p_{\text{Sob}}$ and, in particular, for $N = 1$, i.e., for (1.1). The point is that, in the natural rescaled sense, such patterns on smaller compact subsets around the point $\{0, 0^-\}$ must be governed by a regular stationary solution W satisfying

$$(5.1) \quad -\Delta^2 W + |W|^{p-1}W = 0, \quad W(0) = 1.$$

In the subcritical range, it is guaranteed that radial solutions of (5.1) are highly oscillatory [26], which makes very unlikely to match such a behaviour with the outer region to get an acceptable blow-up pattern of (1.1) in the $\{x, t, u\}$ -variables. We again refer to [18, § 5], and must admit that a rigorous proof of such a nonexistence is absent.

REFERENCES

- [1] M.M. Ad’jutov and L.A. Lepin, Absence of blowing up similarity structures in a medium with a source for constant thermal conductivity, *Differ. Equat.*, **20** (1984), 1279–1281.
- [2] G. Barbatis, *Explicit estimates on the fundamental solution of higher-order parabolic equations with measurable coefficients*, *J. Differ. Equat.*, **174** (2001), 442–463.
- [3] G. Barbatis, *Sharp heat-kernel estimates for higher-order operators with singular coefficients*, *Proc. Edinb. Math. Soc.* (2), **47** (2004), 53–67.
- [4] J. Bebernes and D. Eberly, *Mathematical Problems in Combustion Theory*, *Appl. Math. Sci.*, Vol. **83**, Springer-Verlag, Berlin, 1989.

- [5] C.J. Budd, V.A. Galaktionov, and J.F. Williams, *Self-similar blow-up in higher-order semilinear parabolic equations*, SIAM J. Appl. Math., **64** (2004), 1775–1809.
- [6] C.J. Chapman and M.R.E. Proctor, *Nonlinear Rayleigh-Benard convection between poorly conducting boundaries*, J. Fluid Mech., **101** (1980), 759–782.
- [7] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill Book Company, Inc., New York/London, 1955.
- [8] Yu.V. Egorov, V.A. Galaktionov, V.A. Kondratiev, and S.I. Pohozaev, *Global solutions of higher-order semilinear parabolic equations in the supercritical range*, Adv. Differ. Equat., **9** (2004), 1009–1038.
- [9] J.D. Evans, V.A. Galaktionov, and J.R. King, *Blow-up similarity solutions of the fourth-order unstable thin film equation*, Euro J. Appl. Math., **18** (2007), 195–231.
- [10] J.D. Evans, V.A. Galaktionov, and J.F. Williams, *Blow-up and global asymptotics of the limit unstable Cahn-Hilliard equation*, SIAM J. Math. Anal., **38** (2006), 64–102.
- [11] D.A. Frank-Kamenetskii, *Towards temperature distributions in a reaction vessel and the stationary theory of thermal explosion*, Doklady Acad. Nauk SSSR **18** (1938), 411–412.
- [12] D.A. Frank-Kamenetskii, *Diffusion and Heat Transfer in Chemical Kinetics*, Plenum Press, New York, 1969.
- [13] V.A. Galaktionov, *On a spectrum of blow-up patterns for a higher-order semilinear parabolic equations*, Proc. Royal Soc. London A, **457** (2001), 1–21.
- [14] V.A. Galaktionov, *Critical global asymptotics in higher-order semilinear parabolic equations*, Int. J. Math. Math. Sci., **60** (2003), 3809–3825.
- [15] V.A. Galaktionov, *Evolution completeness of separable solutions of non-linear diffusion equations in bounded domains*, Math. Meth. Appl. Sci., **27** (2004), 1755–1770.
- [16] V.A. Galaktionov, *Geometric Sturmian Theory of Nonlinear Parabolic Equations and Applications*, Chapman & Hall/CRC, Boca Raton, Florida, 2004.
- [17] V.A. Galaktionov, *On blow-up space jets for the Navier-Stokes equations in $\mathbb{R}^3$ with convergence to Euler equations*, J. Math. Phys., **49** (2008), 113101.
- [18] V.A. Galaktionov, *Five types of blow-up in a semilinear fourth-order reaction-diffusion equation: an analytic-numerical approach*, Nonlinearity, submitted (arXiv:0901.4307).
- [19] V.A. Galaktionov and P.J. Harwin, *Non-uniqueness and global similarity solutions for a higher-order semilinear parabolic equation*, Nonlinearity, **18** (2005), 717–746.
- [20] V.A. Galaktionov and S.A. Posashkov, *Application of a new comparison theorem for unbounded solutions of nonlinear parabolic equations*, Differ. Equat., **22** (1986), 809–815.
- [21] V.A. Galaktionov and A.E. Shishkov, *Higher-order quasilinear parabolic equations with singular initial data*, Comm. Contemp. Math., **8** (2006), 331–354.
- [22] V.A. Galaktionov and S.R. Svirshchevskii, *Exact Solutions and Invariant Subspaces of Nonlinear Partial Differential Equations in Mechanics and Physics*, Chapman & Hall/CRC, Boca Raton, Florida, 2007.
- [23] V.A. Galaktionov and J.L. Vazquez, *Continuation of blow-up solutions of nonlinear heat equations in several space dimensions*, Comm. Pure Appl. Math., **50** (1997), 1–68.
- [24] V.A. Galaktionov and J.L. Vazquez, *A Stability Technique for Evolution Partial Differential Equations. A Dynamical Systems Approach*, Progr. in Nonl. Differ. Equat. and Their Appl., Vol. **56**, Birkhäuser, Boston/Berlin, 2004.
- [25] V.A. Galaktionov and J.F. Williams, *Blow-up in a fourth-order semilinear parabolic equation from explosion-convection theory*, Euro J. Appl. Math., **14** (2003), 745–764.

- [26] F. Gazzola and H.-C. Grunau, *Radial entire solutions for supercritical biharmonic equations*, Math. Ann., **334** (2006), 905–936.
- [27] V.L. Gertsberg and G.I. Sivashinsky, *Large cells in nonlinear rayleigh-benard convection*, Prog. Theor. Phys., **66** (1981), 1219–1229.
- [28] Y. Giga and R.V. Kohn, *Asymptotically self-similar blow-up of semilinear heat equations*, Comm. Pure Appl. Math., **38** (1985), 297–319.
- [29] R. Hamilton, *The formation of singularities in the Ricci flow. Surveys in Differ. Geom.*, Vol. II (Cambridge, MA, 1993), pp. 7-136, Int. Press, Cambridge, MA, 1995.
- [30] G. Joulin, A.B. Mikishev, and G.I. Sivashinsky, *A Semenov-Rayleigh-Benard problem*, Preprint.
- [31] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math., **63** (1934), 193–248.
- [32] A. Lunardi, *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, Birkhäuser, Basel/Berlin, 1995.
- [33] E. Mitidieri and S.I. Pohozaev, *Apriori Estimates and Blow-up of Solutions to Nonlinear Partial Differential Equations and Inequalities*, Proc. Steklov Inst. Math., Vol. **234**, Intern. Acad. Publ. Comp. Nauka/Interperiodica, Moscow, 2001.
- [34] C.V. Pao, *Nonlinear Parabolic and Elliptic Equations*, Plenum Press, New York, 1992.
- [35] F. Planchon and P. Raphaël, *Existence and stability of the log-log blow-up dynamics for the L^2 -critical nonlinear Schrödinger equation in a domain*, Ann. Henri Poincaré, **8** (2007), 1177–1219.
- [36] L.A. Peletier and W.C. Troy, *Spatial Patterns. Higher Order Models in Physics and Mechanics*, Birkhäuser, Boston/Berlin, 2001.
- [37] P. Quittner and P. Souplet, *Superlinear Parabolic Problems and Their Equilibria*, Birkhäuser, 2007.
- [38] A.A. Samarskii, V.A. Galaktionov, S.P. Kurdyumov, and A.P. Mikhailov, *Blow-up in Quasilinear Parabolic Equations*, Walter de Gruyter, Berlin/New York, 1995.
- [39] N. Semenov, *Chemical Kinetics and Chain Reaction*, Clarendon Press, Oxford, 1935.
- [40] M.A. Vainberg and V.A. Trenogin, *Theory of Branching of Solutions of Non-Linear Equations*, Noordhoff Int. Publ., Leiden, 1974.
- [41] Ya.B. Zel'dovich, G.I. Barenblatt, V.B. Librovich, and G.M. Makhviladze, *The Mathematical Theory of Combustion and Explosions*, Consultants Bureau [Plenum], New York, 1985.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY, UK
E-mail address: `vag@maths.bath.ac.uk`